



PRODUCT PORTFOLIO

| Poultry Vaccines



va:ino^{va}® Poultry Live Vaccines



■ VAXXON® CLONE	Active immunization against Newcastle Disease.	4
■ VAXXON® LA SOTA	Active immunization against Newcastle Disease.	5
■ VAXXON® B1 HITCHNER	Active immunization against Newcastle Disease.	6
■ VAXXON® IB-H120	Active immunization against Infectious Bronchitis.	7
■ VAXXON® H120-B1 HITCHNER	Active immunization against Infectious Bronchitis and Newcastle Disease.	8
■ VAXXON® H120-LA SOTA	Active immunization against Infectious Bronchitis and Newcastle Disease.	9
■ VAXXON® H120-CLONE	Active immunization against Infectious Bronchitis and Newcastle Disease.	10
■ VAXXON® CHB	Active immunization against Newcastle Disease and Infectious Bronchitis.	11
■ VAXXON® GUMBORO 2	Active immunization against Infectious Bursal Disease (Gumboro).	12
■ VAXXON® GUMBORO 3	Active immunization against Infectious Bursal Disease (Gumboro).	13
■ VAXXON® IBD IMC	Immune Complex Vaccine against Infectious Bursal Disease (Gumboro).	14

■ VAXXON® REO	Active Immunization against viral Arthritis/Tenosynovitis.	15
■ VAXXON® MG LIVE	Active immunisation against loss in egg production due to Mycoplasma gallisepticum infections.	16
■ VAXXON® ILT	Active immunization against Infectious Laryngotracheitis.	17
■ VAXXON® POX	Active immunization against Fowl Pox.	18
■ VAXXON® SHS	Active immunisation against Swollen Head Syndrome (Avian Metapneumovirus).	19
■ VAXXON® AE	Active immunization against Avian Encephalomyelitis.	20
■ VAXXON® MD CVI-N	Vaccine against Marek's Disease.	21
■ VAXXON® MD HVT CVI-N	Vaccine against Marek's Disease.	22
■ VAXXON® COCCIVET	Vaccine against Coccidiosis in broiler chickens.	23
■ VAXXON® COCCIVET R	Vaccine against Coccidiosis in layer and breeder chickens.	24



va:ino^{va}® Poultry Inactivated Vaccines



VAXXON® AVIFLU	Active immunization against Avian Influenza type A.	25
VAXXON® ND FLU	Active immunisation against Newcastle Disease and Avian Influenza A.	26
VAXXON® ND BROILER	Active immunization and reduction of mortality against Newcastle Disease.	27
VAXXON® ND	Booster vaccination for prophylaxis against Newcastle Disease.	28
VAXXON® ND-IB	Booster vaccination against Newcastle disease and Avian Infectious Bronchitis.	29
VAXXON® ND-IBD	Booster vaccination against Newcastle disease and Avian Infectious Bronchitis and to protect progeny against Gumboro disease.	30
VAXXON® ND-IB-IBD	Booster vaccination against Newcastle disease and Avian Infectious Bronchitis and to protect progeny against Gumboro disease.	31
VAXXON® ND-IB-EDS	Active Immunization against Newcastle Disease, Egg Drop Syndrome (EDS'76) and Avian Infectious Bronchitis, to reduce the decline in eggs caused by adenovirus EDS'76.	32

VAXXON® ND-IB3-EDS	Active Immunization against Newcastle Disease, Egg Drop Syndrome (EDS'76) and Avian Infectious Bronchitis, to reduce the decline in eggs caused by adenovirus EDS'76.	33
VAXXON® ND-IB-IBD-REO	Booster vaccination against Newcastle Disease and Avian Infectious Bronchitis and to protect progeny against Gumboro disease and Reovirus.	34
VAXXON® CORYZA 3	Active immunization against Infectious Coryza caused by Avibacterium paragallinarum.	35
VAXXON® MG INAC	Active immunization against Mycoplasma gallisepticum infection.	36
VAXXON® SHS INAC	Active immunisation against Swollen Head Syndrome.	37
VAXXON® REO INAC	Booster vaccination to protect progeny against Reovirus.	38
VAXXON® FC	Active immunization against Fowl Cholera.	39

VAXXON[®] CLONE



Indication

Active immunization against Newcastle Disease.



Active Substance

Live attenuated Newcastle Disease Virus, strain Clone:
 $\geq 10^6$ EID₅₀/dose, in stabilizing solution.



Administration Route

Eye Drop, Spray, Drinking Water.



Poultry Category

Breeders, Layers, Broilers.



Age of Vaccination

Day-Old-Chicks, onwards.



Presentation

Lyophilized, 10x1000, 10x2500 doses.



Shelf life

Packaged product: 24 months after production date.
Reconstituted product: 2 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON® LA SOTA



Indication

Active immunization against Newcastle Disease.



Active Substance

Live attenuated Newcastle Disease Virus, strain LaSota:
 $\geq 10^6$ EID₅₀/dose, in stabilizing solution.



Administration Route

Drinking water.



Poultry Category

Breeders, Layers, Broilers.



Age of Vaccination

As a booster, from 21 days of age, then every 2-3 months.



Presentation

Lyophilized, 10x1000, 10x2500 doses.



Shelf life

Packaged product: 24 months after production date.
Reconstituted product: 2 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON[®] B1 HITCHNER



Indication

Active immunization against Newcastle Disease.



Active Substance

Live attenuated Newcastle Disease virus strain B1 HITCHNER:
≥ 6.0 log₁₀ EID₅₀/dose.



Administration Route

Eye-drop and Spray.



Poultry Category

Breeders, Layers, Broilers.



Age of Vaccination

Day-Old-Chicks, onwards; booster vaccination after 3 weeks.
For laying hens and breeders, a 3rd dose of vaccine at around 10 weeks of age is recommended.



Presentation

Lyophilized, 10x1000, 10x2500 doses.



Shelf life

Packaged product: 24 months after production date.
Reconstituted product: 2 hours.



Storage conditions

Store and transport in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON[®] IB-H120



Indication

Active immunization against Infectious Bronchitis.



Active Substance

Live attenuated Infectious Bronchitis Virus, strain Massachusetts H120: $\geq 10^3$ EID₅₀/dose, in stabilizing solution.



Administration Route

Eye Drop, Spray, Drinking Water.



Poultry Category

Breeders, Layers, Broilers.



Age of Vaccination

Day-Old-Chicks, onwards; booster vaccination after 3 weeks.



Presentation

Lyophilized, 10x1000, 10x2500 doses.



Shelf life

Packaged product: 24 months after production date.
Reconstituted product: 2 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON[®] H120-B1 HITCHNER



Indication

Active immunization against Infectious Bronchitis and Newcastle Disease.



Active Substance

Live attenuated Infectious Bronchitis virus strain H120:
≥ 3.0 log₁₀ EID₅₀/dose
Live attenuated Newcastle Disease virus strain B1 HITCHNER:
≥ 6.0 log₁₀ EID₅₀/dose.



Administration Route

Eye-drop and Spray.



Poultry Category

Breeders, Layers, Broilers.



Age of Vaccination

Day-Old-Chicks, onwards; booster vaccination after 3 weeks.
For laying hens and breeders, a 3rd dose of vaccine at around 10 weeks of age is recommended.



Presentation

Lyophilized, 10x1000, 10x2500 doses.



Shelf life

Packaged product: 24 months after production date.
Reconstituted product: 2 hours.



Storage conditions

Store and transport in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON[®] H120-LA SOTA



Indication

Active immunization against Infectious Bronchitis and Newcastle Disease.



Active Substance

Live attenuated Infectious Bronchitis virus strain H120:
≥ 3.0 log₁₀ EID₅₀/dose
Live attenuated Newcastle Disease virus strain LA SOTA:
≥ 6.0 log₁₀ EID₅₀/dose.



Administration Route

Drinking water.



Poultry Category

Breeders, Layers, Broilers.



Age of Vaccination

From 21 days of age in chickens previously vaccinated.



Presentation

Lyophilized, 10x1000, 10x2500 doses.



Shelf life

Packaged product: 24 months after production date.
Reconstituted product: 2 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON[®] H120-CLONE



Indication

Active immunization against Infectious Bronchitis and Newcastle Disease.



Active Substance

Live attenuated Newcastle Disease Virus, strain Clone:
 $\geq 10^6$ EID₅₀/dose
Live attenuated Infectious Bronchitis Virus, strain Massachusetts H120:
 $\geq 10^3$ EID₅₀/dose, in stabilizing solution.



Administration Route

Eye Drop, Spray, Drinking Water.



Poultry Category

Breeders, Layers, Broilers.



Age of Vaccination

Day-Old-Chicks, onwards; 2nd dose after 3 weeks.



Presentation

Lyophilized, 10x1000, 10x2500 doses.



Shelf life

Packaged product: 24 months after production date.
Reconstituted product: 2 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON[®] CHB



Indication

Active immunization against Newcastle Disease and Infectious Bronchitis.



Active Substance

Live attenuated Newcastle Disease virus strain CLONE:
≥ 6.0 log₁₀ EID₅₀/dose
Live attenuated Infectious Bronchitis virus strain H120:
≥ 3.0 log₁₀ EID₅₀/dose
Live attenuated Infectious Bronchitis virus strain 28/86:
≥ 3.0 log₁₀ EID₅₀/dose.



Administration Route

Eye drop, spray or drinking water.



Poultry Category

Breeders, Layers, Broilers.



Age of Vaccination

At 1st day of age via spray or eye drop; 2nd vaccination 2-3 weeks after first vaccination; booster before laying period.



Presentation

Lyophilized, 10x1000 doses.



Shelf life

Packaged product: 24 months after production date.
Reconstituted product: 2 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON® GUMBORO 2



Indication

Active immunization against Infectious Bursal Disease (Gumboro).



Active Substance

Live intermediated vaccine against Gumboro Disease Virus:
 $\geq 10^3$ EID₅₀/dose, in stabilizing solution.



Administration Route

Drinking Water.



Poultry Category

Breeders, Layers, Broilers.



Age of Vaccination

From 4 days-of-age onward:

1. MDA 300 if possible to evaluate the antibody level;
2. if not possible to evaluate the antibody level -
1st vaccination at 8-10 days of age via drinking water
2nd vaccination at 18-20 days of age via drinking water.



Presentation

Lyophilized, 10x1000, 10x2500 doses.



Shelf life

Packaged product: 24 months after production date.
Reconstituted product: 2 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON® GUMBORO 3



Indication

Active immunization against Infectious Bursal Disease (Gumboro).



Active Substance

Live intermediate plus Gumboro Disease Virus strain Winterfield 2512/90: $\geq 10^{2.7}$ EID₅₀/dose, in stabilizing solution.



Administration Route

Drinking water. In case of emergency eye drop administration at 1st day of age.



Poultry Category

Breeders, Layers, Broilers.



Age of Vaccination

The chickens can be vaccinated starting from the 10th day of age, but specific antibodies are lower than 500 (ELISA). Do not vaccinate chickens with antibody titres higher than 500 or lower than 300. If not possible to evaluate the antibody level - 1st vaccination: 8-10 days of age via drinking water. 2nd vaccination: 18-20 days of age via drinking water. 3rd vaccination (breeders): 28-32 days of age via drinking water.



Presentation

Lyophilized, 10x1000, 10x2500 doses.



Shelf life

Packaged product: 24 months after production date.
Reconstituted product: 2 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON[®] IBD IMC



Indication

Active immunization of one-day-old broilers chicks or 18- to 19-day-old embryonated broiler hatching eggs, with IBD maternal antibodies against Infectious Bursal Disease, to reduce clinical signs of disease caused by very virulent Avian Infectious Bursal Disease (IBD) viruses.



Active Substance

IBD Suspension, strain W2512, sufficient to obtain $\geq 10^{2.5}$ EID₅₀/dose.
AC-IBD Suspension, sufficient to obtain ≥ 0.005 UN/ml.



Administration Route

Subcutaneous or In Ovo.



Poultry Category

Broilers.



Age of Vaccination

At 18-19 days in embryonated eggs or Day-Old-Chicks (SC).



Presentation

Lyophilized, 10x1000, 10x2000, 10x2500, 10x3000, 10x4000, 10x5000, 10x10000 doses.



Shelf life

Packaged product: 24 months after production date.
Reconstituted product: use immediately.



Storage conditions

Store and transport refrigerated (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON[®] REO



Indication

Active Immunization against Arthritis/Tenosynovitis.



Active Substance

Live Attenuated Arthritis Virus, strain S1133: $\geq 10^7$ TCID₅₀/dose, and Diluents.



Administration Route

Subcutaneous.



Poultry Category

Breeders.



Age of Vaccination

In 4-day-old chicks; booster vaccination at 10 weeks of age.



Presentation

Lyophilized, 10x1000.



Shelf life


Packaged product: 24 months after production date.
Reconstituted product: 2 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON[®] MG LIVE



Indication

For active immunisation against loss in egg production due to Mycoplasma gallisepticum infections.



Active Substance

Mycoplasma gallisepticum K5831B-19 strain; $\geq 10^{6.0}$ CFU/dose.



Administration Route

Eye Drop or Spray.



Poultry Category

Layers.



Age of Vaccination

Chickens from 4 weeks of age.



Presentation

Lyophilized, 1x1000, 1x2000 doses; Diluent 1x30ml, 1x60ml.



Shelf life

Packaged product: 36 months after production date.
Reconstituted product: use immediately.



Storage conditions

Store in a refrigerator (≤ 10 °C).
Do not freeze. Protect from light.



VAXXON® ILT



Indication

Active immunization against Infectious Laryngotracheitis.



Active Substance

Live attenuated Laryngotracheitis virus, strain PV/64:
≥ 2.5 log₁₀ EID₅₀/dose.



Administration Route

Eye Drop.



Poultry Category

Breeders, Layers.



Age of Vaccination

At 4 weeks of age; booster dose between 15-20 weeks of age.



Presentation

Lyophilized, 10x1000 doses.



Shelf life

Packaged product: 24 months after production date.
Reconstituted product: 2 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON[®] POX



Indication

Active immunization against Fowl Pox.



Active Substance

Live attenuated Fowl-pox virus strain Brescia P-1:
 $\geq 3.0 \log_{10} \text{TCID}_{50}/\text{dose}$.



Administration Route

Wing web.



Poultry Category

Breeders, Layers.



Age of Vaccination

Breeders: at 5-15 weeks of age; Layers: one month before laying period.



Presentation

Lyophilized, 1x1000 doses, 10x1000 doses.



Shelf life

Packaged product: 24 months after production date.
Reconstituted product: 2 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON[®] SHS



Indication

Active immunisation against Swollen Head Syndrome (Avian Metapneumovirus).



Active Substance

Live attenuated Swollen Head Syndrome virus strain 1439:
 $\geq 10^{2.3}$ TCID₅₀/dose.



Administration Route

Spray or Eye Drop.



Poultry Category

Breeders, Layers.



Age of Vaccination

From 7 days of age onwards.



Presentation

Lyophilized, 10x1000 doses.



Shelf life

Packaged product: 24 months after production date.
Reconstituted product: 2 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON[®] AE



Indication

Active immunization against Avian Encephalomyelitis.



Active Substance

Live attenuated Avian Encephalomyelitis Virus, strain Calnek 1143:
 $\geq 10^3$ EID₅₀/dose.



Administration Route

Drinking water.



Poultry Category

Layers.



Age of Vaccination

Between 10 and 15 weeks of age.



Presentation

Lyophilized, 10x1000 doses.



Shelf life


Packaged product: 24 months after production date.
Reconstituted product: 3 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON[®] MD CVI-N



Indication

Prevention of Marek's Disease.



Active Substance

Live attenuated chicken Marek's Disease virus strain CVI988-N, at least 2000 CFU/dose, and diluents.



Administration Route

Subcutaneous or In Ovo.



Poultry Category

Breeders, Layers, Broilers.



Age of Vaccination

At 18-19 days in embryonated eggs or Day-Old-Chicks (SC).



Presentation

Frozen, 1000, 2000 doses, including diluent 200ml, 400ml.



Shelf life

Packaged product: 24 months after production date.




Storage conditions

Store and transport frozen in liquid nitrogen at -196 °C. Protect from light.



VAXXON[®] MD HVT CVI-N



Indication

Prevention of Marek's Disease.



Active Substance

Live turkey herpesvirus strain FC-126, at least 2000 CFU/dose;
Live attenuated chicken Marek's Disease virus strain CVI988-N,
at least 2000 CFU/dose, and diluents.



Administration Route

Subcutaneous or In Ovo.



Poultry Category

Breeders, Layers, Broilers.



Age of Vaccination

At 18-19 days in embryonated eggs or Day-Old-Chicks (SC).



Presentation

Frozen, 1000, 2000 doses, including diluent 200ml, 400ml.



Shelf life

Packaged product: 24 months after production date.



Storage conditions

Store and transport frozen in liquid nitrogen at -196 °C.
Protect from light.



VAXXON[®] COCCIVET



Indication

Polyvalent live vaccine against Coccidiosis in broiler chickens.



Active Substance

Live attenuated polyvalent vaccine
Eimeria tenella: 0.1 to 0.6 x 10⁶ sporulated oocysts
Eimeria acervuline: 0.1 to 0.6 x 10⁶ sporulated oocysts
Eimeria máxima 52: 0.6 to 2.6 x 10⁵ sporulated oocysts
Eimeria máxima 47: 0.6 to 2.6 x 10⁵ sporulated oocysts
Eimeria máxima 159: 0.6 to 2.6 x 10⁵ sporulated oocysts
Eimeria mitis: 0.1 to 0.5 x 10⁶ sporulated oocysts
Eimeria praecox: 0.1 to 0.5 x 10⁶ sporulated oocysts.



Administration Route

Spray.



Poultry Category

Broiler chickens.



Age of Vaccination

Day-Old-Chicks.



Presentation

Liquid form; 10x1000 (10ml), 10x2000 (15ml), 10x5000 (20ml), 10x10000 (30ml) doses.



Shelf life

Packaged product: 12 months after production date.
After opening: 2 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C).
Do not freeze. Protect from light.

VAXXON[®] COCCIVET R



Indication

Polyvalent live vaccine against Coccidiosis in layer and breeder chickens.



Active Substance

Live attenuated polyvalent vaccine
Eimeria tenella strain BV 25: 1.0 to 6.0 x 10⁵ sporulated oocysts
Eimeria acervulina strain BV 45: 1.0 to 6.0 x 10⁵ sporulated oocysts
Eimeria maxima strain BV 47: 1.0 to 8.0 x 10⁵ sporulated oocysts
Eimeria maxima strain BV 52: 1.0 to 8.0 x 10⁵ sporulated oocysts
Eimeria brunetti strain BV 300: 2.0 to 7.0 x 10⁵ sporulated oocysts
Eimeria necatrix strain BV 302: 1.0 to 6.0 x 10⁵ sporulated oocysts
Eimeria mitis strain BV 44: 1.0 to 5.0 x 10⁵ sporulated oocysts
Eimeria praecox strain BV 41: 1.0 to 5.0 x 10⁵ sporulated oocysts.



Administration Route

Eye-drop and spray.



Poultry Category

Layer and breeder chickens.



Age of Vaccination

Day-Old-Chicks, onwards.



Presentation

Liquid form; 10x1000 (30ml) doses.



Shelf life

Packaged product: 12 months after production date.
After opening: 2 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Protect from light.



VAXXON[®] AVIFLU



Indication

Active immunization against Avian Influenza type A.



Active Substance

Inactivated Avian Influenza virus A, subtype H9N2 :
≥ 5.0 log₂ HI units/dose.



Administration Route

Broiler: 0.2 ml, subcutaneous
Layers & Breeders: 0.2 ml, revaccination 0.5 ml, subcutaneous or intramuscular.



Poultry Category

Breeders, Layers, Broilers.



Age of Vaccination

Broilers at 1st day of age.
Layers & Breeders from 1st day of age up to 3 weeks of age.
Revaccination at 8 weeks of age and between 12-16 weeks of age recommended.



Presentation

Oil emulsion, 500 ml (2500 doses, revaccination 1000 doses).



Shelf life

Packaged product: 24 months. After opening: 10 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Protect from light.

VAXXON[®] ND FLU



Indication

Active immunisation against Newcastle disease and Avian Influenza A.



Active Substance

Inactivated Newcastle Disease virus, strain Ulster:
≥ 4.0 log₂ HI units
Inactivated Avian Influenza A virus, subtype H9N2:
≥ 5.0 log₂ HI units.



Administration Route

Broiler: 0.2 ml, subcutaneous
Layers & Breeders: 0.2 ml, revaccination 0.5 ml, subcutaneous or intramuscular.



Poultry Category

Breeders, Layers, Broilers.



Age of Vaccination

Broilers at 1st day of age.
Layers & Breeders from 1st day of age up to 3 weeks of age.
Revaccination at 8 weeks of age and between 12-16 weeks of age recommended.



Presentation

Oil emulsion, 500 ml (2500 doses, revaccination 1000 doses).



Shelf life

Packaged product: 24 months. After opening: 10 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Protect from light.

VAXXON[®] ND BROILER



Indication

Active immunization and reduction of mortality against Newcastle Disease.



Active Substance

Inactivated Newcastle Disease virus, strain Ulster:
≥ 4.0 log₂ HI units.



Administration Route

0.1 ml, intramuscular.



Poultry Category

Broilers.



Age of Vaccination

From 1st day of age onwards.



Presentation

Oil emulsion, 250 ml (2500 doses), 500 ml (5000 doses).



Shelf life

Packaged product: 24 months. After opening: 10 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Protect from light.

VAXXON[®] ND



Indication

Booster vaccination for prophylaxis against Newcastle Disease.



Active Substance

Inactivated Newcastle Disease virus, strain Ulster:
≥ 4.0 log₂ HI units/dose.



Administration Route

0.5 ml, intramuscular or subcutaneous.



Poultry Category

Breeders, Layers.



Age of Vaccination

From 14 to 20 weeks of age, before the laying period.



Presentation

Oil emulsion, 500 ml (1000 doses).



Shelf life

Packaged product: 24 months. After opening: 10 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Protect from light.



VAXXON[®] ND-IB



Indication

Booster vaccination against Newcastle Disease and Avian Infectious Bronchitis.



Active Substance

Inactivated Newcastle Disease virus, strain Ulster:
≥ 4.0 log₂ HI units
Inactivated Infectious Bronchitis virus, strain Massachusetts M41:
≥ 6.0 log₂ HI units.



Administration Route

0.5 ml, intramuscular or subcutaneous.



Poultry Category

Breeders, Layers.



Age of Vaccination

From 14 to 20 weeks of age, before the laying period.



Presentation

Oil emulsion, 500 ml (1000 doses).



Shelf life

Packaged product: 24 months. After opening: 10 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Protect from light.

VAXXON[®] ND-IBD



Indication

Booster vaccination of breeders and layers against Newcastle Disease and Avian Infectious Bronchitis and to protect progeny against Gumboro.



Active Substance

Inactivated Newcastle Disease virus, strain Ulster:
≥ 4.0 log₂ HI units
Inactivated IBD virus strain Winterfield 2512:
S/P ≥ 0.25



Administration Route

0.5 ml, intramuscular or subcutaneous.



Poultry Category

Breeders, Layers.



Age of Vaccination

From 14 to 20 weeks of age, before the laying period.



Presentation

Oil emulsion, 500 ml (1000 doses).



Shelf life

Packaged product: 24 months. After opening: 10 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Protect from light.



VAXXON[®] ND-IB-IBD



Indication

Booster vaccination against Newcastle Disease and Avian Infectious Bronchitis and to protect progeny against Gumboro Disease.



Active Substance

Inactivated Newcastle Disease virus, strain Ulster:
≥ 4.0 log₂ HI units
Inactivated Infectious Bronchitis virus, strain Massachusetts M41:
≥ 6.0 log₂ HI units
Inactivated IBD virus strain Winterfield 2512: S/P ≥ 0.25.



Administration Route

0.5 ml, intramuscular or subcutaneous.



Poultry Category

Breeders, Layers.



Age of Vaccination

From 14 to 20 weeks of age, before the laying period.



Presentation

Oil emulsion, 500 ml (1000 doses).



Shelf life

Packaged product: 24 months. After opening: 10 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Protect from light.

VAXXON[®] ND-IB-EDS



Indication

Active Immunization against Newcastle Disease, Egg Drop Syndrome (EDS'76) and Avian Infectious Bronchitis, to reduce the decline in eggs caused by adenovirus EDS'76.



Active Substance

Inactivated Newcastle Disease virus, strain Ulster:
≥ 4.0 log₂ HI units
Inactivated Infectious Bronchitis virus, strain Massachusetts M41:
≥ 6.0 log₂ HI units
Inactivated Egg Drop Syndrome'76, strain Mc Ferran 127:
≥ 7.0 log₂ HI units.



Administration Route

0.5 ml, intramuscular or subcutaneous.



Poultry Category

Breeders, Layers.



Age of Vaccination

At 18-20 weeks of age, before the laying period.



Presentation

Oil emulsion, 500 ml (1000 doses).



Shelf life

Packaged product: 24 months. After opening: 10 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Protect from light.

VAXXON[®] ND-IB3-EDS



Indication

Active Immunization against Newcastle Disease, Egg Drop Syndrome (EDS'76) and Avian Infectious Bronchitis, to reduce the decline in eggs caused by adenovirus EDS'76.



Active Substance

Inactivated Newcastle Disease virus, strain Ulster:
≥ 4.0 log₂ HI units
Inactivated Infectious Bronchitis virus, strain Massachusetts M41:
≥ 6.0 log₂ HI units
Inactivated Infectious Bronchitis virus, strain D274:
≥ 4.0 log₂ HI units
Inactivated Infectious Bronchitis virus, strain D1466:
≥ 4.0 log₂ HI units
Inactivated Egg Drop Syndrome'76, strain Mc Ferran 127:
≥ 7.0 log₂ HI units.



Administration Route

0.5 ml, intramuscular or subcutaneous.



Poultry Category

Breeders, Layers.



Age of Vaccination

At 18-20 weeks of age, before the laying period.



Presentation

Oil emulsion, 500 ml (1000 doses).



Shelf life

Packaged product: 24 months. After opening: 10 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Protect from light.

VAXXON[®] ND-IB-IBD-REO



Indication

Booster vaccination against Newcastle Disease and Avian Infectious Bronchitis and to protect progeny against Gumboro disease and Reovirus.



Active Substance

Inactivated Newcastle Disease virus, strain Ulster:
≥ 4.0 log₂ HI units
Inactivated Infectious Bronchitis virus, strain Massachusetts M41:
≥ 6.0 log₂ HI units
Inactivated IBD virus strain Winterfield 2512:
S/P ≥ 0.25
Inactivated virus of the avian Arthritis / tenosynovitis Avian S1133:
S/P ≥ 0.20.



Administration Route

0.5 ml, intramuscular or subcutaneous.



Poultry Category

Breeders, Layers.



Age of Vaccination

From 14 to 20 weeks of age, before the laying period.



Presentation

Oil emulsion, 500 ml (1000 doses).



Shelf life

Packaged product: 24 months. After opening: 10 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Protect from light.

VAXXON[®] CORYZA 3



Indication

Active immunization against Infectious Coryza caused by *Avibacterium paragallinarum*.



Active Substance

Inactivated cultures of *Avibacterium paragallinarum* serotype A, strain 0083: $\geq 2.3 \log_2$ HI
Inactivated cultures of *Avibacterium paragallinarum* serotype B, strain 0222: $\geq 2.3 \log_2$ SAT
Inactivated cultures of *Avibacterium paragallinarum* serotype C, strain Modesto: $\geq 2.3 \log_2$ SAT.



Administration Route

0.5 ml, intramuscular or subcutaneous.



Poultry Category

Breeders, Layers.



Age of Vaccination

At 6 weeks of age, booster dose at 16 weeks before the laying period.



Presentation

Oil emulsion, 500 ml (1000 doses).



Shelf life

Packaged product: 24 months. After opening: 10 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze. Protect from light.



VAXXON[®] MG INAC



Indication

Active immunization against Mycoplasma gallisepticum infection.



Active Substance

Inactivated Mycoplasma gallisepticum strain S-6:
≥ 6.3 log₂ HI.



Administration Route

0.5 ml, subcutaneous.



Poultry Category

Breeders, Layers.



Age of Vaccination

At about 12 weeks of age; booster at 18-20 weeks of age, before the laying period.



Presentation

Oil emulsion, 500 ml (1000 doses).



Shelf life

Packaged product: 24 months. After opening: 10 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Protect from light.



VAXXON[®] SHS INAC



Indication

Active immunisation against Swollen Head Syndrome.



Active Substance

Inactivated Swollen Head Syndrome virus, strain 1439:
 $\geq 5.0 \log_{10}$ TCID₅₀/dose.



Administration Route

0.5 ml, subcutaneous.



Poultry Category

Breeders, Layers.



Age of Vaccination

At 18 weeks of age, before the laying period.



Presentation

Oil emulsion, 500 ml (1000 doses).



Shelf life

Packaged product: 24 months. After opening: 10 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Protect from light.

VAXXON[®] REO INAC



Indication

Booster vaccination to protect progeny against Reovirus.



Active Substance

Inactivated virus of the avian Arthritis / tenosynovitis Avian S1133:
S/P \geq 0.20.



Administration Route

0.5 ml, intramuscular or subcutaneous.



Poultry Category

Breeders, Layers.



Age of Vaccination

From 14 to 20 weeks of age, before the laying period.



Presentation

Oil emulsion, 500 ml (1000 doses).



Shelf life

Packaged product: 24 months. After opening: 10 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Protect from light.



VAXXON[®] FC



Indication

Active immunization against Fowl Cholera.



Active Substance

Inactivated *Pasteurella multocida* serotype 1: $\geq 7.3 \log_2$ SAT
Inactivated *Pasteurella multocida* serotype 3: $\geq 7.3 \log_2$ SAT
Inactivated *Pasteurella multocida* serotype 4: $\geq 7.3 \log_2$ SAT.



Administration Route

0.5 ml, subcutaneous.



Poultry Category

Breeders, Layers.



Age of Vaccination

At 3-4 weeks of age; booster 4 weeks thereafter.



Presentation

Oil emulsion, 500 ml (1000 doses).



Shelf life

Packaged product: 24 months. After opening: 10 hours.



Storage conditions

Store in a refrigerator (2 °C – 8 °C). Do not freeze.
Protect from light.



vaxxino^{va}®

Disclaimer: This brochure may contain information or may show or refer to products and/or services which are only available in certain countries. A reference to any product/service in this brochure does not imply that such product or such product with the same specifications et cetera, is or will be available at your location. Any product shown or referred to in this brochure may be subject to different regulatory requirements depending on the country of use. Please contact your local different regulatory requirements depending on the country of use. Please contact your local Vaxxinova representative for more information.

Vaxxinova® International BV does not give any warranty or other assurance as to the content of this brochure, its accuracy, completeness, timelessness or fitness for any particular purpose. To the full extent permissible by law, Vaxxinova International BV disclaims all responsibility for any damages or losses (including without limitation, financial loss, damages for loss in business projects, loss of profits or other consequential losses) arising in contract, tort or otherwise from the use of inability to use any material of Vaxxinova, or from any action or decision taken as a result of using Vaxxinova's materials. Vaxxinova disclaims all responsibility and liability (including for negligence) in relation to information mentioned in Vaxxinova's materials. This document or its content is not to be copied or further distributed without the consent of Vaxxinova International BV.

VAXXINOVA, VAXXON and GOVAXX are registered trademarks of Vaxxinova International B.V. (the Netherlands).

Vaxxinova® International BV
Transistorweg 5
6534 AT Nijmegen
The Netherlands
www.vaxxinova.com